

THE



CLEAR

APPROACH TO HEMOSTASIS

- SPECIALLY DESIGNED for the esthetic zone
- TRANSPARENT GEL stays on the sulcus
- LEAVES NO RESIDUE and rinses off easily
- DOES NOT INTERFERE with bonding

ACHIEVE HEMOSTASIS . . . ESTHETICALLY

This viscous yet spreadable 25% aluminum chloride gel quickly stops minor bleeding and sulcular fluid without the mess of a liquid or paste.



Subgingival preparation and bleeding sulcus.



Rub hemostatic firmly against bleeding tissues. Achieve the most rapid hemostasis with the Metal Dento-Infusor™ tip. The clear gel allows easy visibility and rinses away quickly.



Press Ultrapak™ cord into gel in sulcus around tooth preparations. Leave 4–5 minutes.



Remove cord. Firm air/water spray. Air dry. Rub gel into the sulcus again. Leave 1 minute.



Finished restoration 2 weeks post-op. Facilitates great control in the esthetic zone with no residue.

- **SPECIALLY DESIGNED** for the esthetic zone
- **TRANSPARENT GEL** stays on the sulcus
- **LEAVES NO RESIDUE** and rinses off easily
- **DOES NOT INTERFERE** with bonding*



*Data on file



6407 - ViscoStat Clear Dento-Infusor IndiSpense Syringe Kit

1 x 30 ml IndiSpense syringe
20 x Metal Dento-Infusor tips
20 x 1.2 ml empty syringes



6409 - ViscoStat Clear Dento-Infusor Syringe Kit

4 x 1.2 ml syringes
20 x Metal Dento-Infusor tips



6408 - ViscoStat Clear IndiSpense Syringe 1pk

30 ml syringe



6410 - ViscoStat Clear Syringe 20pk

1.2 ml syringes

